

**NPEP CHECKLIST**  
**Bendigo Emergency Department**  
**Non-occupational HIV Exposures**

**1. RISK ASSESSMENT**

- Type of exposure
- Date and time of exposure (NPEP must be started within 72 hours)
- HIV status of source/partner
- Determine risk of HIV and need for NPEP using Tables 2 and 3

**2. NPEP PRESCRIPTION**

- For complex cases call The Alfred Hospital 9076 2000 and speak to the ID registrar
- Dispense 7 day NPEP starter pack of either 2 or 3 drugs
  - ✓ 2 drug NPEP = Tenofovir 300mg/ Emtricitabine 200mg Daily
  - ✓ 3 drug NPEP = Tenofovir 300mg/ Emtricitabine 200mg + Dolutegravir 50mg Daily

**3. BASELINE TESTS**

- Serology for HIV/Hepatitis A,B,C/ Syphilis (*only for sexual exposures*)
- U&E
- STI screen for chlamydia and gonorrhoea (NAAT/PCR)
  - ✓ First pass urine/throat swab/rectal swab (MSM)

**4. MANAGEMENT OF OTHER INFECTIONS**

- **Hepatitis B**
  - ✓ Hx of vaccination = no action
  - ✓ Unvaccinated = give 1<sup>st</sup> dose of HBV vaccine
  - ✓ HBIG **only** if unvaccinated **and** source known HBsAg positive
- **Sexually Transmitted Infections (STI)**
  - ✓ If patient has STI symptoms, treat empirically and collect urine/swabs for chlamydia and gonorrhoea PCR (and viral PCR swab of ulcer, if present)

**5. FOLLOW UP ARRANGEMENTS**

- Fax a follow-up referral to Bendigo Sexual Health Clinic (fax 5441 4200)
- Advise patient to make appointment at Bendigo SHC within 7 days to obtain remaining 21 days of NPEP course

**6. COMPLETE INITIAL CONSULTATION FORM AND FAX VICTORIAN NPEP SERVICE**